

What do we learn from Resonance Production in Heavy Ion Collisions?

Christina Markert §

Physics Department, Yale University, New Haven, CT 06520, USA

Abstract. Resonances with their short life time and strong coupling to the dense and hot medium are suggested as a signature of the early stage of the fireball created in a heavy ion collision [1, 2, 3]. The comparison of resonances with different lifetimes and quark contents may give information about time evolution and density and temperature of during the expanding of fireball medium. Resonances in elementary reactions have been measured since 1960. Resonance production in elementary collisions compared with heavy ion collisions where we expect to create a hot and dense medium may show the direct of influence of the medium on the resonances. This paper shows a selection of the recent resonance measurements from SPS and RHIC heavy ion colliders.

1. Resonances in Medium

In a heavy ion collision an extended hot and dense fireball medium is created. During the expansion of the fireball two freeze-out conditions are defined, chemical and thermal, representing the end of the inelastic and elastic interactions. In a dynamical evolving system produced resonances decay and may get regenerated. Decay daughters of resonances which decay inside the medium may also scatter with other particles from the medium, mostly pions for SPS and RHIC energies. This results in a non-reconstructable resonance from the decay daughters measured in the detector, because the invariant mass of the decay daughters no longer matches that of the parent. The reconstructed resonance signal from the scattered decay daughters is a few hundred MeV broad in the UrQMD model [4, 5] and is therefor not distinguishable from the background distribution. The rescattering and regeneration (pseudo-elastic) process for resonances and their decay particles depend on the individual cross sections and are dominant after chemical but before the kinetic freeze-out. These interactions can result in changes of the reconstructed resonance yields, momentum spectra and widths. Rescattering will decrease the measured resonance yields while the mechanism of regeneration will increase the them. Microscopic model calculations include every step in a heavy ion interaction in terms of elastic and inelastic interactions of hadrons and strings [4, 5]. They are therefore able to describe the rescattering and regeneration contribution on the resonances in a fireball. The prediction of this model (UrQMD) is a signal loss for some of the resonances

§ (Christina.markert@yale.edu)

due to more rescattering than regeneration in the low momentum region $p_T < 1$ GeV for the hadronic decays. The leptonic decay products are not significantly affected by rescattering due to their low cross section with hadrons. Comparisons between the yield and momentum spectra of the hadronic and leptonic decay channels can indicate the magnitude of the rescattering and regeneration contribution between chemical and thermal freeze-out.

2. Resonances in elementary collisions

A resonance is a particle with a higher mass than the corresponding ground state particle with the same quark content. It decays strongly and therefore the lifetime, τ , of the resonance is short (within a few fm/c). This causes a wide spread in energy and a natural width, given by $\Gamma = \hbar/\tau$. These broad states with finite Γ and lifetime τ can be formed by collisions between the same particles into which they decay. The resonances can have hadronic and leptonic decay channels. Since they have a short lifetime they can only be measured by reconstruction using the decay particles measured in the detector. At the beginning of the 1960's the first resonance particle was discovered in a bubble chamber experiment at Berkeley using a kaon beam hitting a proton target. The $K(892)^-$ resonance is formed by a $K^- + p \rightarrow K(892)^- + p$ reaction [6]. The reconstruction of the $K(892)^-$ was done via invariant mass calculation using the decay products of a $K(892)^- \rightarrow K^0 + \pi^-$ decay (see Fig 1). The signal in the invariant mass spectrum peaks around the mass of 890 MeV/ c^2 for the $K^-(892)$ particle. In 1968 Luis Walter Alvarez received the Nobel Prize for resonance particles discovered in 1960.




Figure 1. First discovery of resonance in invariant mass reconstruction spectrum of $K(892)^-$ in K^0 and pion channel [6].

Later experiments observed more resonances, and using an energy scan of the kaon beam measured the total (elastic and inelastic) cross section of the $K^- + p$ scattering. The scattering amplitudes versus the kaon beam momentum for different decay channels of the $\Lambda(1520)$ resonance are shown in Fig. 2. At the kaon beam momentum of 395 MeV/c the cross section peaks in the kinematic region of the $\Lambda(1520)$ resonance.

Theoretical in-medium calculations made using the relativistic chiral SU(3) Lagrangian to describe the resonances by using hadrons instead of quarks and gluons are in good agreement with the rescattering amplitude in Fig. 2 [3]. The solid line is the contribution of the s-, p-, d-waves functions while the dashed line shows only the contribution from the s-wave. The mass and width predictions for resonances within this model will be discussed later in this paper.




Figure 2. First discovery of resonance in invariant mass reconstruction spectrum of $K(892)^-$ in K^0 and pion channel [7].

3. Resonance Reconstruction

The resonances are reconstructed from their observed decay daughters. The Λ s from a $\Sigma(1385)$ and $\Xi(1530)$ decay are reconstructed via topological analysis (see table 1). The resonance signal is obtained by the invariant mass reconstruction of each daughter combination and subtraction of the combinatorial background calculated by mixed event or like-signed techniques. The resonance ratios, spectra and yields are measured at mid-rapidity for RHIC at $\sqrt{s_{NN}} = 200$ GeV and over 4π for SPS at $\sqrt{s_{NN}} = 17$ GeV. The central trigger selection for Au+Au collisions at RHIC takes the 5% or 10% and for Pb+Pb collisions at SPS the 5% of the most central inelastic interactions. The setup for the proton+proton interaction is a minimum bias trigger.

Particle	mass [MeV/ c^2]	width [MeV/ c^2]	lifetime [fm/c]	decay channel
$\Delta(1232)$	1232 ± 2	120 ± 5	1.6	$p + \pi$
$K(892)$	896.1 ± 0.27	50.7 ± 0.6	3.89	$K + \pi$
$\Sigma(1385)$	1385 ± 3	37 ± 2	5.2	$\Lambda(\rightarrow p+\pi) + \pi$
$\Lambda(1520)$	1519.5 ± 1.0	15.6 ± 1.0	12.6	$p + K$
$\phi(1020)$	1019.417 ± 0.014	4.458 ± 0.032	44.6	$K^+ + K^-$

Table 1. Resonances and their main hadronic decay channels from PDG [8]

4. Rescattering and Regeneration

Rescattering will decrease the resonance signal in the invariant mass reconstruction while the regeneration process will increase the signal. If the signal loss or gain is due to rescattering or regeneration a comparison of the resonance yields and momentum spectra with thermal models is not appropriate as the contribution of the rescattering phase has taken to be into account. Microscopic models (such as UrQMD) are able to describe such a phase [4, 5]. Depending on the lifetime and the rescattering and regeneration cross sections the observed resonances come from different times of the fireball source.


**Figure 3.** In UrQMD produced (dashed line) and reconstructable (solid line) resonances versus the fireball time for $\rho(770)$, $\Delta(1232)$, $K(892)$, $\Sigma(1385)$, $\Lambda(1520)$ and $\phi(1020)$ at mid-rapidity for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

Fig. 3 shows the UrQMD produced (dashed line) and reconstructable (solid line) resonances versus the fireball time for different resonances with different lifetimes: $\tau_\rho = 1.3$ fm/c, $\tau_\Delta(1232) = 1.7$ fm/c, $\tau_{K(892)} = 4.0$ fm/c, $\tau_{\Sigma(1385)} = 5.5$ fm/c, $\tau_{\Lambda(1520)} = 13$ fm/c and $\tau_\phi = 46$ fm/c. We observe two features: the mean of the fireball lifetime at the decay of the reconstructable resonances scales with the lifetime of the resonance and observed resonances compared to those produced is higher for states with less rescattering and regeneration. Note that this ratio is not directly applicable to the total yield of produced resonances from a thermal model prediction because it is a dynamical process where resonances decay and get generated several times during the whole expansion time which makes this ratio very small. However, the absolute values of the measurable resonances of UrQMD can be compared with the experimental results. One short remark here is that UrQMD has a long lifetime for a Au+Au heavy ion reaction (longer than 30 fm/c). It is not clear how the resonance production in terms of rescattering and regeneration would be affected if the source would expand faster.

From UrQMD calculations we learn that the signal loss of resonances due to rescattering takes place predominantly in the low momentum region [4, 5]. Therefore the reconstructed transverse momentum spectra are expected to change to higher inverse slope parameters and higher $\langle p_T \rangle$. An exponential fit to the transverse momentum of the created and measurable resonances calculated in UrQMD shows that the spectra shape looks thermal before and after rescattering and regeneration of the resonance Fig 4. The increase in $\langle p_T \rangle$ is for the $\Delta(1232)$ and $K(892)$ is of the order of 100 MeV/c and for the $\phi(1020)$ 35 MeV/c.




Figure 4. Transverse momentum spectrum of produced and reconstructable resonances in UrQMD $\Delta(1232)$, $K(892)$ and $\phi(1020)$ at mid-rapidity for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

We also would expect a higher increase of inverse slope and $\langle p_T \rangle$ for resonances from p+p to Au+Au collisions than for ground state particles due to the rescattering and regeneration effect in the surrounding fireball medium even in peripheral collisions. The STAR data from p+p and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV confirm this trend of a strong increase $\langle p_T \rangle$ for resonances from p+p to peripheral Au+Au collisions which is

not present for the ground state particles (see Fig 5) [9, 10, 11]. Due to this observation one would also expect a stronger deviation of the transverse momentum spectra of the resonances compared to ground state particles in the low momentum region of a thermal model.




Figure 5. The $\langle p_T \rangle$ for resonances and ground state particles in p+p and Au+Au collisions versus number of charged particles [9, 10, 11, 19].

5. Resonance Yields

Under the assumption that all the particles freeze out at the same time their yields are extracted from a thermal model for one fit parameter set of chemical temperature T_{ch} , chemical potentials μ_b and μ_s and the strangeness saturation factor γ_s . This chemical freeze-out is the end of the inelastic interactions where the yields of most particle species are constant. Elastic and pseudo-elastic interactions do not change the yield of the ground state particles. The momentum spectra can change due to further elastic interactions. Resonances, which are measured (reconstructed) by their decay daughters, are affected by the pseudo-elastic interactions which can cause a change in the yield of the measured signal in the invariant mass spectrum and a change in the momentum distribution of the reconstructed resonance. The yields and momentum spectra will change for resonances according to their cross section for rescattering and regeneration of resonances and rescattering of their decay daughters, the resonance lifetime and the medium density. Regeneration may compensate the effects of rescattering. Yields and momentum spectra in comparison with the prediction from a thermal model may suggest the magnitude of influence of rescattering over regeneration. Resonance over

non-resonance particle ratios of heavy ion collisions compared to p+p interactions may indicate signal yield changes due to rescattering and regeneration processes. Table 2 shows measured particle ratios in p+p and Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV and the expected particle ratios for the resonances from a thermal model based on a fit to the ground state particle ratios for Au+Au collisions [12].

Particle ratio	p+p data	Au+Au data	Au+Au model
$\Delta(1232)/p$	0.72 ± 0.108	0.96 ± 0.148	0.68
$K(892)^0/K$	0.389 ± 0.029	0.228 ± 0.044	0.32
$\Lambda(1520)/\Lambda$	0.104 ± 0.03	0.033 ± 0.017	0.071
$\phi(1020)/K$	0.124 ± 0.025	0.1556 ± 0.0311	0.13

Table 2. Comparison of STAR data with thermal model predictions of resonance/non-resonance ratios in p+p and Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV [12].

The $K(892)/K$ and the $\Lambda(1520)/\Lambda$ in Au+Au collisions are lower than in p+p collisions and the expected values from the thermal model are also higher than the data (see Fig 7). These data suggest that the rescattering cross section is larger than the regeneration from the $K+\pi$ and $K+p$. The $K(892)$ lifetime is smaller than the lifetime of the $\Lambda(1520)$ which would imply a larger suppression for $K(892)/K$ than for the $\Lambda(1520)/\Lambda$ ratio. The competing contribution of regeneration seems to be larger for $K(892)$ than for the $\Lambda(1520)$. The $\phi(1020)/K$ ratio is consistent within errors with the thermal model prediction, which is expected because only a small fraction of the $\phi(1020)$ are decaying inside the fireball due to the long lifetime of the $\phi(1020)$, of 46 fm/c. The expected contribution of rescattering for the short lived $\Delta(1232)$ resonance is larger than that for the $K(892)$ and the $\Lambda(1520)$. However the $\Delta(1232)/p$ ratio does not decrease from p+p to Au+Au collisions and is on the order of $41\% \pm 22\%$ higher than the thermal model prediction. This indicates a large cross section for the regeneration of $\Delta(1232)$ resonance in the $p+\pi$ channel. The first $\Sigma(1385)$ yields from heavy ion collisions appear to follow the same trend as the $\Delta(1232)$ [13]. This implies that the $\Lambda+\pi$ regeneration cross section is compensating the signal loss from rescattering. From this observation we can conclude that there is a ranking order of the cross section for the different regeneration processes: $\sigma_{p+\pi} \geq \sigma_{\Lambda+\pi} > \sigma_{K+\pi} > \sigma_{K+p}$. The first attempt to measure the $\Xi(1530)$ with the STAR detector are shown in [14]. Final results will give contributions to the $\Xi(1530) + \pi$ cross section and add a more stringent test to theoretical descriptions of the data.

6. Time Scale

Depending on the length of the time interval between chemical and kinetic freeze-out, $\Delta\tau$, the magnitude of the suppression factor of the measured resonance will change due to contributions from rescattering and regeneration. A model using thermally produced particle yields at chemical freeze-out and an additional rescattering phase, including the lifetime of the resonances and decay product interactions within the expanding fireball of nuclear matter, can place a lower limit on [15, 16, 17]. The probability of rescattering is described by the (energy averaged) interaction cross section of the decay particles of the resonances within the medium, and depends on the radius, density and velocity of the fireball. A slower expansion of the fireball would lead to a higher suppression of the resonance signal due to the larger rate of rescattering. This model does not include regeneration and is therefore predicting a lower limit of the lifetime $\Delta\tau$ between the two freeze-out surfaces. It is only applicable if the contribution from rescattering is larger than from regeneration. The two ratios $K(892)/K$ and $\Lambda(1520)/\Lambda$ are expected to have a larger rescattering contribution. A $\Delta\tau > 4$ fm/c results if chemical freeze-out occurs at 160 MeV (See Fig 6.




Figure 6. Dependence of lifetime and chemical freeze out temperature of a fireball source given by the two particle ratios $\Lambda(1520)/\Lambda$ and $K(892)/K$ including limits from the STAR data for central Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [15, 16].

This model gives every combination between temperature and life span in the marked area of the data and their errors. Another valid interpretation of the data in this model is a chemical freeze-out temperature of $T=110 \pm 10$ MeV and a time span of $\Delta\tau = 0$ fm/c. With smaller errors the combination of T and $\Delta\tau$ can be determined

more precisely. Fig. 7 shows the resonance/non-resonance ratios in p+p and Au+Au collisions for different centralities. The suppression of the $K(892)/K$ and the $\Lambda(1520)/\Lambda$ in Au+Au collisions sets in at the peripheral collisions and remains constant up to the central collisions. This would suggest the same life span between chemical and thermal freeze-out in mid-peripheral and central Au+Au collisions.




Figure 7. Resonance/non-resonance ratios of $\phi(1020)/K^-$ [9], $\Delta(1232)^{++}/p$, $\rho(770)/\pi$ [18], $K(892)/K^-$ [11] and $\Lambda(1520)/\Lambda$ [20, 21] for p+p and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The ratios are normalized to the $K(892)/K^-$ p+p ratio. Statistical and systematical errors are included.

7. Leptonic and Hadronic Decay Channels

Direct comparisons of the spectra and yields in heavy ion collisions from leptonic and hadronic decay channels may show the influence of the hadronic interaction phase after chemical freeze-out folded with the initial production of the $\phi(1020)$ at the early stage. The $\phi(1020)$ is one of the resonances where we have measurements from the leptonic and hadronic decay channels in heavy ion collisions. At the moment there are only SPS data from heavy ion collisions available for the leptonic channel of $\phi(1020)$. Results from $\phi(1020)$ production in Au+Au collisions at RHIC energies into $e^+ + e^-$ should be measured in the next months. From SPS we have the so-called ' ϕ puzzle' with two different momentum distributions and yields for the leptonic and hadronic decay. Fig 8 shows the transverse momentum distribution of the hadronic decay $\phi \rightarrow K^+ + K^-$ from NA49 and the leptonic decay $\phi \rightarrow \mu^+ + \mu^-$ from NA50 [22, 23]. The inverse slope parameter from fits to the momentum spectra indicated as lines are $T = 305 \pm 15$ MeV

for hadronic decay and $T = 218 \pm 10$ MeV for leptonic decay. The extracted yield from the momentum spectrum of the leptonic decay is a factor of 4 ± 2 higher than the one for the hadronic decay. Microscopic calculations (UrQMD) estimated for the $\phi(1020)$ resonance a suppression of 20-30% of the yield in the hadronic decay channel due to rescattering of the kaon decay daughters in the low momentum region $p_T < 1$ GeV [4, 5]. The rescattering is negligible for the leptonic decay due to the very low cross section of interaction with the hadronic phase. The trend of the lower signal in the low momentum region of the hadronic decay compared to the leptonic decay from the data is in agreement with the model. However this signal loss of 20-30% from the model calculation is not sufficient to explain the factor of 4 ± 2 in the measured yield of the data.




Figure 8. Transverse momentum distribution of the hadronic decay $\phi(1020) \rightarrow K^+ + K^-$ from NA49 [22] and the leptonic decay $\phi(1020) \rightarrow \mu^+ + \mu^-$ from NA50 [23].

This gives room for possible medium effects on the resonance production which is effective at an earlier stage, before chemical freeze-out. An approach to describe the in medium modification of the $\phi(1020)$ resonance has been done by K. Haglin and E. Kolomeitsev [24, 25]. This attempt to fit the SPS data describes a hot and dense fireball where the lifetime of the $\phi(1020)$ resonance is modified towards smaller lifetimes and therefore more of the $\phi(1020)$ resonances decay inside the medium. This will introduce a larger signal loss due to rescattering of the hadronic decay daughters. In this conference the $\phi(1020)$ measurements from PHENIX in d+Au [26] and in Au+Au [27] and in d+Au collision systems were shown.

8. Width and Mass

Theoretical in-medium calculations of resonances predict mass and width changes depending on the nuclear matter density. The theory of Hendrik van Hees and Ralf Rapp has predictions for modifications to the $\Delta(1232)^{++}$ width at chemical and kinetic freeze-out temperatures of $T_{\text{ch}} = 180$ MeV $T_{\text{kn}} = 120$ MeV, which corresponds to densities of $\rho_{\text{ch}} = 0.68 \rho_0$ and $\rho_{\text{kn}} = 0.12 \rho_0$ [28]. Calculations from M. Lutz [2, 3] predict a mass shift and widths broadening of 40 MeV and 100 MeV for the $\Sigma(1385)$ and the $\Lambda(1520)$ at a medium density of $\rho = 1 \rho_0$. Additional calculations which include the density evolution of the fireball and further rescattering and regeneration processes of the resonances need to be done to give final answer on the measured mass position and width. The measured data from SPS and RHIC do not show a mass shift or width broadening in the errors of 5 MeV. At the moment we don't know if this is a result that we would expect from theoretical calculations.

References

- [1] R. Rapp and J. Wambach, *Adv. Nucl. Phys.* **25** (2000) 1.
- [2] M.F.M. Lutz and C.L. Korpa, *nucl-th/0105067*.
- [3] M.F.M. Lutz, *J.Phys.* **G28** (2002) 1729.
- [4] M. Bleicher and J. Aichelin, *Phys. Lett.* **B530** (2002) 81, *hep-ph/0201123*.
- [5] M. Bleicher private communication.
- [6] M. Alstone et al., *Phys. Rev. Lett.* **6** (1961) 300.
- [7] T.S. Mast et al., *Phys. Rev.* **D14** (1976) 14.
- [8] Particle Data Group, *Eur. Phys. J.* **C3** (1998).
- [9] J. Ma et al., *J.Phys.* **G30** (2004) 543.
- [10] C. Markert et al., *J.Phys.* **G30** (2004) 1313.
- [11] H. Zhang et al., *J.Phys.* **G30** (2004) 577.
- [12] P. Braun-Munzinger, D. Magestro, K. Redlich and J. Stachel, *Phys. Lett.* **B518** (2001) 41, private communication.
- [13] S. Salur, this Proceedings.
- [14] R. Witt, this Proceedings.
- [15] G. Torrieri and J. Rafelski, *Phys. Lett.* **B509** (2001) 239-245.
- [16] G. Torrieri and J. Rafelski, *J.Phys.* **G28** (2002) 1911-1920.
- [17] C. Markert, G. Torrieri, J. Rafelski, Campos do Jordao 2002, New states of matter in hadronic interactions 533-552, *hep-ph/0206260*.
- [18] P. Fachini et al., *J.Phys.* **G30** (2004) 565.
- [19] S. Salur et al., International School of Subnuclear Physics, Erice, Italy 29 August- 7 September 2004 (Submitted to *EPJ C*).
- [20] L. Gaudichet *J.Phys.* **G30** (2004) 549.
- [21] C. Markert et al., *Proc. 19th Winter Workshop on Nuclear Dynamics* (2003) 71.
- [22] V. Friese et al., *J.Phys.* **G23** (1997) 1837.
- [23] N. Willis et al., *Nucl. Phys.* **A661** (1999) 534.
- [24] K. Haglin, *Nucl. Phys.* **A584** (1995) 719, K. Haglin this Proceedings.
- [25] E. Kolomeitsev and P. Filip, *J. Phys.* **G28** (2002) 1697.
- [26] D. Pal, this Proceedings.
- [27] D. Mukhopadhyay, this Proceedings.
- [28] H. van Hees, R. Rapp, this Proceedings.